



## NQF - Female Measures

E & M Codes: Clinic Only

Encounter Date(s): Jan 1, 2009 through Dec 31, 2009

Location	Provider	Breast Cancer Screening	Cervical Cancer Screening	Chlamydia Screening	Osteoporosis Management in Women who had a Fracture
SETMA 1	Aziz	50.3%	0.3%	--	20.0%
	Duncan	66.9%	11.8%	48.0%	50.0%
	Henderson	60.6%	12.3%	26.3%	25.0%
	Murphy	44.1%	2.5%	0.0%	40.0%
	Thomas	47.0%	0.2%	0.0%	--
<b>SETMA 1 Totals:</b>		56.6%	7.5%	35.4%	31.2%
SETMA 2	Anthony	66.6%	0.3%	0.0%	20.0%
	Anwar	76.6%	0.3%	0.0%	0.0%
	Colbert	--	--	0.0%	--
	Cricchio, M	66.0%	0.4%	33.3%	0.0%
	Holly	80.3%	0.0%	0.0%	0.0%
	Leifeste	80.4%	5.1%	0.0%	--
	Wheeler	72.0%	21.7%	1.9%	0.0%
<b>SETMA 2 Totals:</b>		73.3%	8.9%	3.0%	3.7%
SETMA West	Curry	59.0%	13.5%	36.8%	25.0%
	Halbert	60.0%	3.1%	50.0%	0.0%
	Horn	75.9%	8.7%	63.0%	--
	Satterwhite	70.7%	12.4%	46.7%	0.0%
	Vardiman	53.6%	0.0%	0.0%	--
	Young	53.9%	2.8%	0.0%	0.0%
<b>SETMA West Totals:</b>		65.7%	8.4%	48.5%	10.0%
<b>SETMA Totals:</b>		65.3%	8.3%	29.4%	13.2%

Breast Cancer Screening - Yearly mammography screening for women 40 to 69 years of age.

Cervical Cancer Screening - Cervical cancer screening for women 21 to 64 years of age at least every two years.

Chlamydia Screening - Sexually active females between the ages of 16 and 24 should have a Chlamydia screening at least yearly.



## **NQF - Female Measures**

---

E & M Codes: Clinic Only

Encounter Date(s): Jan 1, 2009 through Dec 31, 2009

---

Osteoporosis Management In Women Who Had A Fracture - Women 67 years of age and older who suffered a fracture and who had either (1) a bone mineral density test or (2) a prescription for a drug to treat osteoporosis within six months following the fracture.