



## NQF - Pediatric Measures

E & M Codes: Clinic Only  
 Encounter Date(s): Jan 1, 2009 through Dec 31, 2009

Location	Provider	Childhood Immunization Status	Appropriate Treatment for Children with Pharyngitis
SETMA 1	Aziz	--	0.0%
	Duncan	--	10.0%
	Henderson	0.0%	22.2%
	Murphy	--	0.0%
	Thomas	--	33.3%
<b>SETMA 1 Totals:</b>		0.0%	15.6%
SETMA 2	Anthony	6.1%	17.1%
	Anwar	--	--
	Colbert	3.2%	40.7%
	Cricchio, M	6.2%	37.5%
	Holly	--	--
	Leifeste	--	0.0%
	Wheeler	0.0%	53.5%
<b>SETMA 2 Totals:</b>		3.6%	36.7%
SETMA West	Anthony	--	--
	Curry	8.3%	50.0%
	Halbert	0.0%	0.0%
	Horn	0.0%	69.7%
	Satterwhite	7.7%	69.2%
	Vardiman	--	0.0%
	Young	--	16.7%
<b>SETMA West Totals:</b>		5.6%	43.8%
<b>SETMA Totals:</b>		3.6%	36.2%

Childhood Immunizations - Children should have the following immunizations by their second birthday: four diphtheria, tetanus and acellular pertussis (DTaP/DT), three polio (IPV), one measles, mumps and rubella (MMR), two H influenza type B (HiB) three hepatitis B, one chicken pox (VZV) and four pneumococcal conjugate vaccines.

Appropriate Testing for Children with Pharyngitis - Patients 2 to 18 years of age who were diagnosed with Pharyngitis and given an antibiotic and a strep test for the episode.