



HEDIS - Effectiveness of Acute Care

E & M Codes: Clinic Only
 Encounter Date(s): Jan 1, 2010 through Dec 31, 2010

Location	Provider	Avoidance of Antibiotic Treatment in Adults with Acute Bronchitis	Appropriate Testing for Children with Pharyngitis	Appropriate Treatment for Children with Upper Respiratory Infection
SETMA 1	Anthony	--	--	--
	Aziz	30.8%	--	--
	Duncan	19.6%	6.2%	0.0%
	Henderson	33.3%	30.3%	15.4%
	Murphy	35.7%	0.0%	0.0%
	Thomas	16.4%	100.0%	--
	SETMA 1 Totals:		21.5%	23.5%
SETMA 2	Anthony	10.4%	39.4%	0.0%
	Anwar	100.0%	--	--
	Colbert	--	54.2%	15.7%
	Cricchio, A	--	--	--
	Cricchio, M	84.4%	57.0%	50.0%
	Holly	--	--	--
	Leifeste	50.0%	--	--
	Wheeler	5.6%	56.3%	30.0%
SETMA 2 Totals:		36.1%	52.1%	15.4%
SETMA West	Curry	12.5%	66.7%	6.2%
	Deiparine	21.4%	0.0%	--
	Halbert	11.8%	0.0%	--
	Horn	12.0%	69.0%	0.0%
	Qureshi	12.5%	50.0%	0.0%
	Satterwhite	40.0%	100.0%	0.0%
	Vardiman	38.0%	25.0%	--
	Young	32.7%	33.3%	0.0%
SETMA West Totals:		22.7%	43.1%	2.5%
SETMA Totals:		24.1%	48.7%	12.8%



HEDIS - Effectiveness of Acute Care

E & M Codes: Clinic Only

Encounter Date(s): Jan 1, 2010 through Dec 31, 2010

Avoidance of Antibiotic Treatment in Adults with Acute Bronchitis - Patients 18 to 64 years of age who were diagnosed with acute bronchitis and not prescribed an antibiotic.

Appropriate Testing for Children with Pharyngitis - Patients 2 to 18 years of age who were diagnosed with Pharyngitis and given an antibiotic and a strep test for the episode.

Appropriate Treatment for Children with Upper Respiratory Infection - Patients 3 months to 18 years of age who were diagnosed with upper respiratory infection and not prescribed an antibiotic.