



NQF - Care for Older Adults

E & M Codes: Clinic Only
 Encounter Date(s): Jan 1, 2010 Through Dec 30, 2010

Location	Provider	Counseling for Physical Activity	Colorectal Cancer Screening	Fall Risk Assessment	Urinary Incontinence Assessment
SETMA 1	Anthony	--	--	--	--
	Aziz	46.3%	62.7%	99.3%	79.6%
	Duncan	87.7%	68.7%	96.8%	92.0%
	Henderson	50.3%	68.6%	100.0%	99.2%
	Murphy	60.4%	55.5%	98.3%	88.8%
	Thomas	41.8%	46.6%	97.3%	96.5%
SETMA 1 Totals:		56.6%	60.4%	98.5%	89.9%
SETMA 2	Anthony	99.0%	73.4%	99.9%	96.3%
	Anwar	98.2%	81.5%	99.6%	91.4%
	Colbert	--	--	--	--
	Cricchio, A	100.0%	100.0%	100.0%	100.0%
	Cricchio, M	97.6%	74.7%	99.7%	99.4%
	Holly	98.2%	71.5%	99.5%	98.6%
	Leifeste	88.0%	81.4%	94.3%	96.9%
	Wheeler	98.5%	70.8%	99.3%	97.3%
SETMA 2 Totals:		96.7%	76.7%	98.8%	95.5%
SETMA West	Curry	98.0%	60.6%	99.9%	85.1%
	Deiparine	92.4%	45.9%	99.3%	92.1%
	Halbert	68.4%	50.9%	98.8%	95.1%
	Horn	69.9%	68.8%	100.0%	98.2%
	Qureshi	52.6%	64.3%	100.0%	97.0%
	Satterwhite	85.1%	65.4%	99.2%	95.0%
	Vardiman	72.2%	60.0%	98.9%	84.7%
	Young	91.6%	50.4%	100.0%	97.1%
SETMA West Totals:		78.8%	57.2%	99.3%	93.5%
SETMA Totals:		78.0%	64.6%	98.9%	93.1%



NQF - Care for Older Adults

E & M Codes: Clinic Only

Encounter Date(s): Jan 1, 2010 Through Dec 30, 2010

Counseling For Physical Activity - Patients 65 and older should be counseled to engage in physical activity.

Colorectal Cancer Screening - Patients 50 to 80 years of age should have at least one of the following: a colonoscopy within the last nine years, a flexible sigmoidoscopy within the last four years, a double contrast barium enema within the last four years or a fecal occult blood test within the last year.

Fall Risk Assessment - Patients 65 and older should have a fall risk assessment completed at least yearly.

Urinary Incontinence Assessment - Patients 65 and older should have urinary incontinence assessed at least every six months.