



## NQF - Female Measures

E & M Codes: Clinic Only

Encounter Date(s): Jan 1, 2010 through Dec 31, 2010

Location	Provider	Breast Cancer Screening	Cervical Cancer Screening	Chlamydia Screening	Osteoporosis Management in Women who had a Fracture
SETMA 1	Aziz	43.4%	34.7%	--	0.0%
	Duncan	47.0%	48.1%	35.4%	100.0%
	Henderson	53.6%	49.8%	16.1%	33.3%
	Murphy	47.5%	34.5%	0.0%	--
	Thomas	36.4%	24.9%	0.0%	0.0%
<b>SETMA 1 Totals:</b>		47.1%	42.2%	25.6%	22.2%
SETMA 2	Anthony	54.2%	33.7%	11.1%	0.0%
	Anwar	63.8%	29.3%	--	0.0%
	Cricchio, M	50.3%	23.6%	0.0%	0.0%
	Holly	55.3%	42.5%	--	0.0%
	Leifeste	64.5%	40.5%	0.0%	66.7%
	Wheeler	40.1%	53.0%	3.6%	0.0%
<b>SETMA 2 Totals:</b>		52.9%	39.7%	4.3%	11.1%
SETMA West	Curry	52.4%	43.8%	100.0%	100.0%
	Deiparine	57.8%	45.0%	0.0%	--
	Halbert	47.4%	25.8%	0.0%	0.0%
	Horn	58.5%	36.9%	46.7%	--
	Qureshi	47.1%	49.6%	33.3%	--
	Satterwhite	53.9%	36.3%	0.0%	--
	Vardiman	50.9%	33.6%	0.0%	0.0%
	Young	56.6%	20.3%	33.3%	0.0%
<b>SETMA West Totals:</b>		54.7%	36.5%	38.3%	16.7%
<b>SETMA Totals:</b>		52.1%	38.9%	22.3%	15.2%

Breast Cancer Screening - Yearly mammography screening for women 40 to 69 years of age.

Cervical Cancer Screening - Cervical cancer screening for women 21 to 64 years of age at least every two years.



## **NQF - Female Measures**

---

E & M Codes: Clinic Only

Encounter Date(s): Jan 1, 2010 through Dec 31, 2010

---

Chlamydia Screening - Sexually active females between the ages of 16 and 24 should have a Chlamydia screening at least yearly.

Osteoporosis Management In Women Who Had A Fracture - Women 67 years of age and older who suffered a fracture and who had either (1) a bone mineral density test or (2) a prescription for a drug to treat osteoporosis within six months following the fracture.