



NQF - Pediatric Measures

E & M Codes: Clinic Only
 Encounter Date(s): Jan 1, 2010 through Dec 31, 2010

Location	Provider	Childhood Immunization Status	Appropriate Treatment for Children with Pharyngitis
SETMA 1	Anthony	--	--
	Aziz	0.0%	--
	Duncan	--	6.2%
	Henderson	33.3%	30.3%
	Murphy	0.0%	0.0%
	Thomas	--	100.0%
SETMA 1 Totals:		25.0%	23.5%
SETMA 2	Anthony	4.2%	39.4%
	Anwar	--	--
	Colbert	2.3%	54.2%
	Cricchio, A	--	--
	Cricchio, M	--	57.0%
	Holly	--	--
	Leifeste	--	--
	Wheeler	1.7%	56.3%
SETMA 2 Totals:		2.5%	52.1%
SETMA West	Curry	0.0%	66.7%
	Deiparine	--	0.0%
	Halbert	--	0.0%
	Horn	0.0%	69.0%
	Qureshi	--	50.0%
	Satterwhite	0.0%	100.0%
	Vardiman	--	25.0%
	Young	0.0%	33.3%
SETMA West Totals:		0.0%	43.1%
SETMA Totals:		2.7%	48.7%



NQF - Pediatric Measures

E & M Codes: Clinic Only
Encounter Date(s): Jan 1, 2010 through Dec 31, 2010

Childhood Immunizations - Children should have the following immunizations by their second birthday: four diphtheria, tetanus and acellular pertussis (DTaP/DT), three polio (IPV), one measles, mumps and rubella (MMR), two H influenza type B (HiB) three hepatitis B, one chicken pox (VZV) and four pneumococcal conjugate vaccines.

Appropriate Testing for Children with Pharyngitis - Patients 2 to 18 years of age who were diagnosed with Pharyngitis and given an antibiotic and a strep test for the episode.