



HEDIS - Effectiveness of Acute Care

E & M Codes: Clinic Only
 Encounter Date(s): Jan 1, 2011 through Dec 31, 2011

Location	Provider	Avoidance of Antibiotic Treatment in Adults with Acute Bronchitis	Appropriate Testing for Children with Pharyngitis	Appropriate Treatment for Children with Upper Respiratory Infection
SETMA 1	Aziz	31.8%	--	100.0%
	Duncan	15.3%	22.2%	0.0%
	Halbert	--	--	--
	Henderson	15.7%	38.7%	12.5%
	Murphy	60.0%	0.0%	0.0%
	Palang	14.3%	--	--
	Thomas	19.2%	0.0%	--
SETMA 1 Totals:		19.2%	31.4%	10.0%
SETMA 2	Abbas	--	--	--
	Anthony	5.0%	33.3%	0.0%
	Anwar	75.0%	--	--
	Colbert	--	60.8%	15.3%
	Cricchio, A	0.0%	100.0%	0.0%
	Cricchio, M	71.7%	59.5%	14.3%
	Deiparine	--	--	--
	Holly	--	--	--
	Leifeste	--	--	--
	Murphy	--	--	--
	Wheeler	33.3%	67.3%	7.7%
SETMA 2 Totals:		42.2%	57.0%	14.1%
SETMA West	Curry	13.3%	37.5%	6.7%
	Deiparine	24.1%	6.7%	5.6%
	Halbert	9.1%	--	0.0%
	Horn	26.3%	78.6%	0.0%
	Qureshi	8.2%	69.2%	14.3%
	Satterwhite	0.0%	0.0%	0.0%



HEDIS - Effectiveness of Acute Care

E & M Codes: Clinic Only
 Encounter Date(s): Jan 1, 2011 through Dec 31, 2011

Location	Provider	Avoidance of Antibiotic Treatment in Adults with Acute Bronchitis	Appropriate Testing for Children with Pharyngitis	Appropriate Treatment for Children with Upper Respiratory Infection
SETMA West	Vardiman	59.0%	20.0%	100.0%
SETMA West Totals:		30.5%	43.5%	7.5%
SETMA Totals:		27.5%	53.0%	12.2%

Avoidance of Antibiotic Treatment in Adults with Acute Bronchitis - Patients 18 to 64 years of age who were diagnosed with acute bronchitis and not prescribed an antibiotic.

Appropriate Testing for Children with Pharyngitis - Patients 2 to 18 years of age who were diagnosed with Pharyngitis and given an antibiotic and a strep test for the episode.

Appropriate Treatment for Children with Upper Respiratory Infection - Patients 3 months to 18 years of age who were diagnosed with upper respiratory infection and not prescribed an antibiotic.