



NQF - Care for Older Adults

E & M Codes: Clinic Only

Encounter Date(s): Jan 1, 2011 Through Dec 30, 2011

Location	Provider	Counseling for Physical Activity	Colorectal Cancer Screening	Fall Risk Assessment	Urinary Incontinence Assessment
SETMA 1	Aziz	52.8%	63.7%	99.0%	95.1%
	Duncan	90.8%	68.2%	94.8%	99.2%
	Halbert	100.0%	--	100.0%	100.0%
	Henderson	53.1%	65.3%	99.8%	99.7%
	Murphy	69.6%	60.6%	98.6%	98.4%
	Palang	40.2%	36.9%	98.0%	97.4%
	Thomas	31.0%	39.8%	90.6%	98.2%
SETMA 1 Totals:		59.9%	59.4%	98.0%	97.8%
SETMA 2	Abbas	0.0%	100.0%	100.0%	100.0%
	Anthony	99.2%	69.5%	99.4%	98.9%
	Anwar	98.6%	78.5%	98.6%	97.2%
	Colbert	--	--	--	--
	Cricchio, A	12.8%	58.0%	85.9%	93.1%
	Cricchio, M	97.7%	72.7%	99.3%	99.9%
	Deiparine	--	0.0%	--	--
	Holly	99.1%	77.4%	100.0%	99.6%
	Leifeste	88.9%	81.8%	97.7%	98.3%
	Murphy	--	100.0%	--	--
	Wheeler	98.9%	74.5%	97.3%	98.7%
SETMA 2 Totals:		91.9%	74.8%	97.8%	98.1%
SETMA West	Curry	86.1%	64.2%	99.8%	89.6%
	Deiparine	52.0%	52.6%	98.6%	98.7%
	Halbert	73.3%	52.4%	98.9%	98.3%
	Horn	94.0%	65.5%	99.7%	99.1%
	Qureshi	47.1%	61.4%	99.6%	97.4%
	Satterwhite	88.1%	56.2%	99.8%	89.9%
	Vardiman	77.0%	56.1%	99.1%	94.2%



NQF - Care for Older Adults

E & M Codes: Clinic Only
Encounter Date(s): Jan 1, 2011 Through Dec 30, 2011

Location	Provider	Counseling for Physical Activity	Colorectal Cancer Screening	Fall Risk Assessment	Urinary Incontinence Assessment
SETMA West Totals:		73.6%	57.7%	99.2%	96.3%
SETMA Totals:		75.1%	63.5%	98.3%	97.4%

Counseling For Physical Activity - Patients 65 and older should be counseled to engage in physical activity.

Colorectal Cancer Screening - Patients 50 to 80 years of age should have at least one of the following: a colonoscopy within the last nine years, a flexible sigmoidoscopy within the last four years, a double contrast barium enema within the last four years or a fecal occult blood test within the last year.

Fall Risk Assessment - Patients 65 and older should have a fall risk assessment completed at least yearly.

Urinary Incontinence Assessment - Patients 65 and older should have urinary incontinence assessed at least every six months.