



## NQF - Female Measures

E & M Codes: Clinic Only

Encounter Date(s): Jan 1, 2012 through Dec 31, 2012

Location	Provider	Breast Cancer Screening	Cervical Cancer Screening	Chlamydia Screening	Osteoporosis Management in Women who had a Fracture
SETMA 1	Aziz	45.7%	66.5%	0.0%	14.3%
	Deiparine	40.0%	59.1%	--	--
	Duncan	47.9%	76.5%	2.0%	0.0%
	Henderson	53.6%	72.1%	3.6%	16.7%
	Holly	60.8%	55.2%	--	0.0%
	Murphy	35.3%	52.7%	0.0%	14.3%
	Palang	43.9%	54.5%	0.0%	40.0%
	Thomas	28.6%	70.4%	0.0%	--
<b>SETMA 1 Totals:</b>		46.2%	67.7%	2.5%	17.2%
SETMA 2	Anthony	57.0%	55.3%	12.5%	0.0%
	Anwar	65.6%	63.3%	0.0%	14.3%
	Cash	44.1%	52.2%	0.0%	--
	Colbert	--	--	0.0%	--
	Holly	0.0%	0.0%	--	--
	LaBorde	50.0%	56.7%	0.0%	--
	Leifeste	74.8%	72.6%	0.0%	100.0%
	Read	48.8%	63.3%	0.0%	0.0%
	Wheeler	49.5%	81.1%	1.5%	0.0%
<b>SETMA 2 Totals:</b>		57.1%	68.8%	2.0%	6.2%
SETMA Mid County	Castro	66.7%	92.3%	--	--
	George	36.4%	42.9%	--	--
	Read	52.1%	60.6%	0.0%	--
	Shepherd	56.4%	78.2%	--	0.0%
	Thomas	34.0%	70.1%	0.0%	--
<b>SETMA Mid County Totals:</b>		49.6%	72.0%	0.0%	0.0%
SETMA West	Darden	52.1%	69.2%	0.0%	--



## NQF - Female Measures

E & M Codes: Clinic Only

Encounter Date(s): Jan 1, 2012 through Dec 31, 2012

Location	Provider	Breast Cancer Screening	Cervical Cancer Screening	Chlamydia Screening	Osteoporosis Management in Women who had a Fracture
SETMA West	Deiparine	52.9%	72.8%	0.0%	--
	Halbert	28.4%	33.5%	0.0%	0.0%
	Horn	45.3%	62.3%	5.9%	20.0%
	Qureshi	36.8%	53.6%	0.0%	--
	Vardiman	31.0%	47.4%	33.3%	0.0%
<b>SETMA West Totals:</b>		43.4%	59.8%	3.7%	5.6%
<b>SETMA Totals:</b>		48.1%	64.8%	2.7%	9.5%

Breast Cancer Screening - Yearly mammography screening for women 40 to 69 years of age.

Cervical Cancer Screening - Cervical cancer screening for women 21 to 64 years of age at least every two years.

Chlamydia Screening - Sexually active females between the ages of 16 and 24 should have a Chlamydia screening at least yearly.

Osteoporosis Management In Women Who Had A Fracture - Women 67 years of age and older who suffered a fracture and who had either (1) a bone mineral density test or (2) a prescription for a drug to treat osteoporosis within six months following the fracture.