



NQF - Pediatric Measures

E & M Codes: Clinic Only

Encounter Date(s): Jan 1, 2012 through Dec 31, 2012

Location	Provider	Childhood Immunization Status	Appropriate Treatment for Children with Pharyngitis
SETMA 1	Aziz	--	100.0%
	Deiparine	--	--
	Duncan	--	54.5%
	Halbert	--	--
	Henderson	44.4%	21.1%
	Holly	--	--
	Murphy	--	25.0%
	Palang	--	--
	Thomas	--	--
SETMA 1 Totals:		44.4%	34.3%
SETMA 2	Anthony	11.2%	21.0%
	Anwar	--	--
	Cash	--	0.0%
	Colbert	39.8%	58.7%
	Holly	--	--
	LaBorde	--	--
	Leifeste	0.0%	0.0%
	Read	--	40.0%
	Wheeler	3.8%	62.3%
SETMA 2 Totals:		33.8%	48.9%
SETMA Mid County	Castro	--	--
	George	--	100.0%
	Read	--	--
	Shepherd	--	0.0%
	Thomas	--	--
SETMA Mid County Totals:		--	25.0%



NQF - Pediatric Measures

E & M Codes: Clinic Only
 Encounter Date(s): Jan 1, 2012 through Dec 31, 2012

Location	Provider	Childhood Immunization Status	Appropriate Treatment for Children with Pharyngitis
SETMA West	Darden	--	31.6%
	Deiparine	--	0.0%
	Halbert	--	0.0%
	Horn	0.0%	11.1%
	Qureshi	--	66.7%
	Vardiman	--	33.3%
SETMA West Totals:		0.0%	24.4%
SETMA Totals:		33.9%	45.3%

Childhood Immunizations - Children should have the following immunizations by their second birthday: four diphtheria, tetanus and acellular pertussis (DTaP/DT), three polio (IPV), one measles, mumps and rubella (MMR), two H influenza type B (HiB) three hepatitis B, one chicken pox (VZV) and four pneumococcal conjugate vaccines.

Appropriate Testing for Children with Pharyngitis - Patients 2 to 18 years of age who were diagnosed with Pharyngitis and given an antibiotic and a strep test for the episode.