



## HEDIS - Effectiveness of Acute Care

E & M Codes: Clinic Only  
 Encounter Date(s): Jan 1, 2013 through Dec 31, 2013

Location	Provider	Avoidance of Antibiotic Treatment in Adults with Acute Bronchitis	Appropriate Testing for Children with Pharyngitis	Appropriate Treatment for Children with Upper Respiratory Infection
SETMA 1	Aziz	0.0%	--	--
	Duncan	10.0%	45.5%	12.5%
	Foster	0.0%	9.1%	57.1%
	Henderson	100.0%	17.6%	5.9%
	Holly	--	--	--
	Le	13.0%	38.7%	12.5%
	Murphy	100.0%	0.0%	--
	Palang	22.2%	--	--
	Shepherd	--	--	--
	Thomas	--	--	--
<b>SETMA 1 Totals:</b>		18.8%	26.8%	23.4%
SETMA 2	Anthony	--	32.7%	0.0%
	Anwar	--	--	--
	Cash	--	--	--
	Colbert	--	52.1%	0.0%
	Foster	--	--	--
	Leifeste	100.0%	0.0%	100.0%
	Read	--	19.0%	--
	Wheeler	100.0%	82.7%	20.0%
<b>SETMA 2 Totals:</b>		100.0%	53.8%	20.0%
SETMA Mid County	Castro	60.0%	0.0%	--
	George	25.0%	--	--
	Read	--	--	--
	Shepherd	40.0%	--	--
	Thomas	0.0%	--	--
<b>SETMA Mid County Totals:</b>		35.0%	0.0%	--



## HEDIS - Effectiveness of Acute Care

E & M Codes: Clinic Only  
 Encounter Date(s): Jan 1, 2013 through Dec 31, 2013

Location	Provider	Avoidance of Antibiotic Treatment in Adults with Acute Bronchitis	Appropriate Testing for Children with Pharyngitis	Appropriate Treatment for Children with Upper Respiratory Infection
SETMA Orange	Anwar	--	--	--
	Aziz	0.0%	--	--
	Castro	--	--	--
	Cox	--	100.0%	0.0%
	Green	0.0%	76.5%	--
	Holly	--	--	--
	Le	--	--	--
	Shepherd	--	--	--
<b>SETMA Orange Totals:</b>		<b>0.0%</b>	<b>81.0%</b>	<b>0.0%</b>
SETMA West	Darden	100.0%	40.0%	--
	Deiparine, C	--	0.0%	100.0%
	Halbert	--	0.0%	--
	Horn	33.3%	70.0%	0.0%
	Qureshi	0.0%	0.0%	0.0%
<b>SETMA West Totals:</b>		<b>11.8%</b>	<b>44.0%</b>	<b>11.1%</b>
<b>SETMA Totals:</b>		<b>22.5%</b>	<b>49.2%</b>	<b>20.8%</b>

Avoidance of Antibiotic Treatment in Adults with Acute Bronchitis - Patients 18 to 64 years of age who were diagnosed with acute bronchitis and not prescribed an antibiotic.

Appropriate Testing for Children with Pharyngitis - Patients 2 to 18 years of age who were diagnosed with Pharyngitis and given an antibiotic and a strep test for the episode.

Appropriate Treatment for Children with Upper Respiratory Infection - Patients 3 months to 18 years of age who were diagnosed with upper respiratory infection and not prescribed an antibiotic.