



NQF - Care for Older Adults

E & M Codes: Clinic Only
 Encounter Date(s): Jan 1, 2013 Through Dec 30, 2013

Location	Provider	Counseling for Physical Activity	Colorectal Cancer Screening	Fall Risk Assessment	Urinary Incontinence Assessment
SETMA 1	Aziz	54.4%	66.6%	96.1%	92.4%
	Duncan	90.3%	68.2%	97.8%	99.8%
	Foster	92.0%	65.0%	99.7%	98.2%
	Henderson	62.3%	66.8%	99.7%	99.8%
	Holly	96.5%	73.9%	99.8%	97.9%
	Le	29.4%	45.3%	99.8%	87.0%
	Murphy	67.0%	62.9%	97.0%	98.9%
	Palang	62.0%	57.8%	99.4%	99.7%
	Thomas	42.9%	41.8%	95.2%	92.9%
SETMA 1 Totals:		65.7%	62.7%	98.2%	97.4%
SETMA 2	Anthony	96.1%	74.6%	99.9%	99.3%
	Anwar	96.4%	79.0%	100.0%	98.2%
	Cash	99.0%	59.7%	99.6%	93.2%
	Colbert	--	--	--	--
	Foster	100.0%	60.3%	100.0%	97.9%
	Leifeste	89.6%	83.5%	98.3%	98.1%
	Read	87.2%	74.4%	95.7%	98.6%
	Wheeler	97.8%	69.6%	99.9%	99.1%
SETMA 2 Totals:		94.4%	73.3%	98.9%	97.7%
SETMA Mid County	Castro	96.6%	58.7%	99.7%	98.1%
	George	57.7%	48.7%	100.0%	96.6%
	Read	100.0%	100.0%	100.0%	100.0%
	Shepherd	84.8%	72.5%	99.9%	96.9%
	Thomas	50.1%	37.0%	98.3%	95.9%
SETMA Mid County Totals:		79.8%	57.6%	99.6%	97.1%
SETMA Orange	Anwar	97.8%	69.0%	100.0%	99.3%
	Aziz	42.4%	53.7%	95.7%	83.7%



NQF - Care for Older Adults

E & M Codes: Clinic Only
 Encounter Date(s): Jan 1, 2013 Through Dec 30, 2013

Location	Provider	Counseling for Physical Activity	Colorectal Cancer Screening	Fall Risk Assessment	Urinary Incontinence Assessment
SETMA Orange	Castro	98.0%	60.7%	100.0%	100.0%
	Cox	61.1%	33.2%	97.9%	25.9%
	Green	60.8%	29.0%	98.2%	48.7%
	Holly	100.0%	65.0%	100.0%	100.0%
	Le	--	50.0%	--	--
	Shepherd	83.1%	66.8%	99.4%	95.3%
SETMA Orange Totals:		71.7%	42.8%	98.5%	61.5%
SETMA West	Darden	51.3%	62.4%	99.4%	99.1%
	Deiparine, C	89.7%	62.3%	96.7%	96.4%
	Halbert	80.1%	54.4%	96.4%	92.2%
	Horn	93.1%	67.1%	99.9%	98.6%
	Qureshi	59.6%	60.7%	99.2%	98.4%
SETMA West Totals:		78.6%	60.8%	97.9%	95.9%
SETMA Totals:		78.7%	63.2%	98.6%	95.6%

Counseling For Physical Activity - Patients 65 and older should be counseled to engage in physical activity.

Colorectal Cancer Screening - Patients 50 to 80 years of age should have at least one of the following: a colonoscopy within the last nine years, a flexible sigmoidoscopy within the last four years, a double contrast barium enema within the last four years or a fecal occult blood test within the last year.

Fall Risk Assessment - Patients 65 and older should have a fall risk assessment completed at least yearly.

Urinary Incontinence Assessment - Patients 65 and older should have urinary incontinence assessed at least every six months.