



NQF - Pediatric Measures

E & M Codes: Clinic Only

Encounter Date(s): Jan 1, 2013 through Dec 31, 2013

Location	Provider	Childhood Immunization Status	Appropriate Treatment for Children with Pharyngitis
SETMA 1	Aziz	--	--
	Duncan	--	45.5%
	Foster	13.5%	9.1%
	Henderson	100.0%	17.6%
	Holly	--	--
	Le	0.0%	38.7%
	Murphy	--	0.0%
	Palang	--	--
	Shepherd	--	--
	Thomas	--	--
SETMA 1 Totals:		9.3%	26.8%
SETMA 2	Anthony	7.8%	32.7%
	Anwar	--	--
	Cash	--	--
	Colbert	22.3%	52.1%
	Foster	--	--
	Leifeste	--	0.0%
	Read	--	19.0%
	Wheeler	10.3%	82.7%
SETMA 2 Totals:		18.4%	53.8%
SETMA Mid County	Castro	--	0.0%
	George	--	--
	Read	--	--
	Shepherd	--	--
	Thomas	--	--
SETMA Mid County Totals:		--	0.0%



NQF - Pediatric Measures

E & M Codes: Clinic Only
 Encounter Date(s): Jan 1, 2013 through Dec 31, 2013

Location	Provider	Childhood Immunization Status	Appropriate Treatment for Children with Pharyngitis
SETMA Orange	Anwar	--	--
	Aziz	--	--
	Castro	--	--
	Cox	--	100.0%
	Green	0.0%	76.5%
	Holly	--	--
	Le	--	--
	Shepherd	--	--
SETMA Orange Totals:		0.0%	81.0%
SETMA West	Darden	--	40.0%
	Deiparine, C	--	0.0%
	Halbert	--	0.0%
	Horn	--	70.0%
	Qureshi	--	0.0%
SETMA West Totals:		--	44.0%
SETMA Totals:		17.4%	49.2%

Childhood Immunizations - Children should have the following immunizations by their second birthday: four diphtheria, tetanus and acellular pertussis (DTaP/DT), three polio (IPV), one measles, mumps and rubella (MMR), two H influenza type B (HiB) three hepatitis B, one chicken pox (VZV) and four pneumococcal conjugate vaccines.

Appropriate Testing for Children with Pharyngitis - Patients 2 to 18 years of age who were diagnosed with Pharyngitis and given an antibiotic and a strep test for the episode.