



## NQF - Pediatric Measures

E & M Codes: Clinic Only  
 Encounter Date(s): Jan 1, 2014 through Dec 31, 2014

Location	Provider	Childhood Immunization Status	Appropriate Treatment for Children with Pharyngitis
SETMA 1	Aziz	--	--
	Duncan	--	100.0%
	Foster	6.0%	53.5%
	Henderson	0.0%	0.0%
	Holly	--	--
	Le	0.0%	75.0%
	McClure	--	--
	Murphy	--	100.0%
	Palang	--	0.0%
	Thomas	--	--
<b>SETMA 1 Totals:</b>		<b>4.6%</b>	<b>56.6%</b>
SETMA 2	Anthony	3.9%	37.5%
	Anwar	--	--
	Cash	--	--
	Colbert	8.3%	45.8%
	Cricchio, M	100.0%	50.0%
	Foster	--	--
	Leifeste	--	--
	Read	--	--
	Smith	9.2%	66.7%
	Wheeler	1.4%	70.9%
<b>SETMA 2 Totals:</b>		<b>6.4%</b>	<b>55.3%</b>
SETMA Lumberton	Anwar	--	--
	Arcala, A	--	--
	Foster	0.0%	--
	Halbert	--	--



## NQF - Pediatric Measures

E & M Codes: Clinic Only  
 Encounter Date(s): Jan 1, 2014 through Dec 31, 2014

Location	Provider	Childhood Immunization Status	Appropriate Treatment for Children with Pharyngitis
SETMA Lumberton	Holly	--	--
	Le	0.0%	54.5%
	Leifeste	--	--
	Read	--	0.0%
	Thomas	--	--
<b>SETMA Lumberton Totals:</b>		0.0%	40.0%
SETMA Mid County	Castro	--	--
	Cricchio, M	--	--
	George	--	--
	Shepherd	--	--
	Thomas	--	--
<b>SETMA Mid County Totals:</b>		--	--
SETMA Orange	Anwar	--	--
	Arcala, A	--	--
	Aziz	--	--
	Castro	--	--
	Cox	--	50.0%
	Green	--	57.9%
	Le	--	--
	Shepherd	--	--
<b>SETMA Orange Totals:</b>		--	56.5%
SETMA PA	Arcala, A	--	--
	Arcala, F	--	--
	Castro	--	--
	Cricchio, M	--	--
	Shepherd	--	--



## NQF - Pediatric Measures

E & M Codes: Clinic Only  
 Encounter Date(s): Jan 1, 2014 through Dec 31, 2014

Location	Provider	Childhood Immunization Status	Appropriate Treatment for Children with Pharyngitis
<b>SETMA PA Totals:</b>		--	--
SETMA West	Cricchio, M	--	--
	Darden	--	0.0%
	Deiparine, C	--	0.0%
	Halbert	--	0.0%
	Horn	--	25.0%
	Qureshi	--	50.0%
<b>SETMA West Totals:</b>		--	16.7%
<b>SETMA Totals:</b>		6.1%	51.9%

Childhood Immunizations - Children should have the following immunizations by their second birthday: four diphtheria, tetanus and acellular pertussis (DTaP/DT), three polio (IPV), one measles, mumps and rubella (MMR), two H influenza type B (HiB) three hepatitis B, one chicken pox (VZV) and four pneumococcal conjugate vaccines.

Appropriate Testing for Children with Pharyngitis - Patients 2 to 18 years of age who were diagnosed with Pharyngitis and given an antibiotic and a strep test for the episode.