



Diabetes Consortium - HgbA1c Measures

E & M Codes: Clinic Only

Encounter Date(s): Jan 1, 2015 through Dec 31, 2015

Report Criteria: Patients 18 to 75 With a Chronic Diagnosis of Diabetes
Specialists Excluded (Dr. Ahmed Included)

| Location | Provider | HgbA1c Level | | | | HgbA1c Frequency | | |
|--------------------------------|-------------|--------------|--------|-------|-------------|------------------|-----------------|---------------------|
| | | <= 6.5 | <= 7.0 | > 7.0 | Not Present | Within 3 Months | Within 6 Months | Not Within 6 Months |
| SETMA 1 | Aziz | 32.5% | 48.6% | 50.5% | 0.6% | 81.6% | 93.9% | 6.1% |
| | Duncan | 38.3% | 53.9% | 45.4% | 0.3% | 64.7% | 82.7% | 17.3% |
| | Henderson | 40.0% | 57.9% | 41.5% | 0.4% | 68.0% | 86.2% | 13.8% |
| | Holly | 48.1% | 64.2% | 34.9% | 0.9% | 82.1% | 93.4% | 6.6% |
| | Le | 34.8% | 57.6% | 41.4% | 1.0% | 63.1% | 83.2% | 16.8% |
| | Murphy | 43.2% | 61.8% | 37.6% | 0.2% | 80.7% | 92.4% | 7.6% |
| | Palang | 39.3% | 56.4% | 42.3% | 1.3% | 70.7% | 83.7% | 16.3% |
| | Thomas | 54.5% | 63.6% | 36.4% | 0.0% | 72.7% | 90.9% | 9.1% |
| SETMA 1 Totals: | | 39.1% | 56.8% | 42.4% | 0.6% | 72.6% | 87.6% | 12.4% |
| SETMA 2 | Akhter | 13.2% | 23.3% | 75.6% | 0.7% | 88.0% | 94.7% | 5.3% |
| | Anthony | 32.2% | 53.8% | 46.2% | 0.0% | 94.7% | 98.2% | 1.8% |
| | Anwar | 39.6% | 53.1% | 46.4% | 0.2% | 80.7% | 93.6% | 6.4% |
| | Cash | 20.5% | 34.5% | 64.4% | 0.1% | 93.4% | 97.5% | 2.5% |
| | Cricchio, M | 25.4% | 38.1% | 23.8% | 38.1% | 38.1% | 50.8% | 49.2% |
| | Kansara | 40.7% | 55.2% | 40.7% | 4.1% | 70.3% | 85.8% | 14.2% |
| | Smith | 30.0% | 48.1% | 37.4% | 14.5% | 54.5% | 73.1% | 26.9% |
| | Wheeler | 47.5% | 57.2% | 42.4% | 0.2% | 80.0% | 91.3% | 8.7% |
| SETMA 2 Totals: | | 27.6% | 41.5% | 56.1% | 1.9% | 84.7% | 92.9% | 7.1% |
| SETMA Lumberton | Anwar | 46.2% | 53.8% | 46.2% | 0.0% | 84.6% | 92.3% | 7.7% |
| | Dao | 41.3% | 57.2% | 42.0% | 0.8% | 69.5% | 88.0% | 12.0% |
| | Elliott | 27.7% | 47.9% | 46.2% | 5.0% | 67.2% | 76.5% | 23.5% |
| | Foster | 25.7% | 42.0% | 56.4% | 1.6% | 61.9% | 80.0% | 20.0% |
| | Le | 42.9% | 66.7% | 33.3% | 0.0% | 38.1% | 81.0% | 19.0% |
| | Read | 34.2% | 49.6% | 48.5% | 1.9% | 61.6% | 81.4% | 18.6% |
| SETMA Lumberton Totals: | | 33.1% | 49.4% | 48.8% | 1.7% | 64.3% | 82.5% | 17.5% |



Diabetes Consortium - HgbA1c Measures

E & M Codes: Clinic Only
 Encounter Date(s): Jan 1, 2015 through Dec 31, 2015

Report Criteria: Patients 18 to 75 With a Chronic Diagnosis of Diabetes
 Specialists Excluded (Dr. Ahmed Included)

| Location | Provider | HgbA1c Level | | | | HgbA1c Frequency | | |
|---------------------------------|--------------|--------------|--------|-------|-------------|------------------|-----------------|---------------------|
| | | <= 6.5 | <= 7.0 | > 7.0 | Not Present | Within 3 Months | Within 6 Months | Not Within 6 Months |
| SETMA Mid County | Arcala, A | 28.9% | 36.8% | 52.6% | 10.5% | 47.4% | 65.8% | 34.2% |
| | Arcala, F | 39.2% | 60.8% | 39.2% | 0.0% | 64.7% | 84.3% | 15.7% |
| | Castro | 37.1% | 57.2% | 42.1% | 0.6% | 86.8% | 94.1% | 5.9% |
| | Cricchio, M | 37.8% | 52.8% | 42.8% | 4.4% | 67.4% | 86.2% | 13.8% |
| | George | 50.9% | 67.4% | 30.3% | 2.3% | 64.6% | 82.9% | 17.1% |
| | Shepherd | 47.6% | 64.7% | 34.0% | 1.3% | 81.2% | 93.3% | 6.7% |
| | Thomas | 40.8% | 51.7% | 46.6% | 1.7% | 77.7% | 90.8% | 9.2% |
| SETMA Mid County Totals: | | 42.1% | 58.6% | 39.4% | 2.0% | 77.2% | 90.2% | 9.8% |
| SETMA Orange | Castro | 36.4% | 47.7% | 52.3% | 0.0% | 95.5% | 100.0% | 0.0% |
| | Cox | 31.7% | 51.0% | 43.3% | 5.7% | 63.0% | 79.4% | 20.6% |
| | Green | 31.9% | 50.7% | 41.7% | 7.6% | 47.8% | 67.0% | 33.0% |
| | Read | 40.9% | 72.7% | 27.3% | 0.0% | 68.2% | 84.1% | 15.9% |
| | Shepherd | 61.1% | 69.4% | 30.6% | 0.0% | 94.4% | 97.2% | 2.8% |
| SETMA Orange Totals: | | 33.3% | 52.2% | 42.0% | 5.9% | 59.2% | 75.9% | 24.1% |
| SETMA PA | Arcala, A | 18.1% | 26.4% | 56.9% | 16.7% | 61.1% | 79.2% | 20.8% |
| | Arcala, F | 24.9% | 35.6% | 58.8% | 5.6% | 71.8% | 91.5% | 8.5% |
| SETMA PA Totals: | | 22.9% | 32.9% | 58.2% | 8.8% | 68.7% | 88.0% | 12.0% |
| SETMA West | Darden | 37.0% | 57.5% | 41.3% | 1.2% | 63.3% | 79.2% | 20.8% |
| | Deiparine, C | 28.1% | 44.3% | 52.7% | 3.0% | 66.3% | 81.0% | 19.0% |
| | Halbert | 42.0% | 60.3% | 37.0% | 2.4% | 57.2% | 72.2% | 27.8% |
| | Horn | 39.7% | 57.7% | 40.9% | 1.2% | 65.3% | 81.0% | 19.0% |
| | Khan | 40.2% | 55.0% | 42.2% | 2.5% | 48.4% | 70.5% | 29.5% |
| | Qureshi | 32.9% | 44.8% | 54.6% | 0.6% | 77.1% | 91.5% | 8.5% |
| SETMA West Totals: | | 36.9% | 53.6% | 44.5% | 1.9% | 63.2% | 79.2% | 20.8% |



Diabetes Consortium - HgbA1c Measures

E & M Codes: Clinic Only

Encounter Date(s): Jan 1, 2015 through Dec 31, 2015

Report Criteria: Patients 18 to 75 With a Chronic Diagnosis of Diabetes
Specialists Excluded (Dr. Ahmed Included)

| Location | Provider | HgbA1c Level | | | | HgbA1c Frequency | | |
|----------------------|----------|--------------|--------|-------|-------------|------------------|-----------------|---------------------|
| | | <= 6.5 | <= 7.0 | > 7.0 | Not Present | Within 3 Months | Within 6 Months | Not Within 6 Months |
| SETMA Totals: | | 34.5% | 50.6% | 47.2% | 2.0% | 73.4% | 86.7% | 13.3% |